



**Injury Risk for Research Subjects With Spina
Bifida Occulta in a
Repeated Impact Study:
A Case Review**

By

**John P. Albano
Samuel G. Shannon
Nabih M. Alem
Kevin T. Mason**

Aircrew Protection Division

19960226 099

January 1996

DTIC QUALITY INSPECTED 1

Approved for public release; distribution unlimited.

**U.S. Army Aeromedical Research Laboratory
Fort Rucker, Alabama 36362-0577**

Notice

Qualified requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

Change of address

Organizations receiving reports from the U.S. Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.


Disposition

Destroy this document when it is no longer needed. Do not return it to the originator.

Disclaimer

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Reviewed:




KEVIN T. MASON
LTC(P), MC, MFS
Director, Aircrew Protection
Division

Released for publication:



FOR ROGER W. WILEY, O.D., Ph.D.
Chairman, Scientific
Review Committee



DENNIS F. SHANAHAN
Colonel, MC, MFS
Commanding

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release, distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) USAARL Report No. 96-05			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Aeromedical Research Laboratory		6b. OFFICE SYMBOL (If applicable) MCMR-UAD	7a. NAME OF MONITORING ORGANIZATION U.S. Army Medical Research and Materiel Command		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 620577 Fort Rucker, AL 36362-0577			7b. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21702-5012		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-91-C-1115		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62787A	PROJECT NO. 30162787A878	TASK NO. FA
11. TITLE (Include Security Classification) Injury risk for research subjects with spina bifida occulta in a repeated impact study: A case review					
12. PERSONAL AUTHOR(S) John P. Albano, Samuel G. Shannon, Nabih M. Alem, and Kevin T. Mason					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day)	
15. PAGE COUNT					
16. SUPPLEMENTAL NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) repeated impact, spina bifida occulta, axial loading		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Spina bifida occulta (SBO) occurs in 18-34% of the normal U. S. population. Recently, 16.5% of normal, asymptomatic male soldier volunteer candidates in a U.S. Army Aeromedical Research Laboratory ride motion study were excluded from the study because they had SBO at one vertebral level. Disqualifying this percentage of screened research subject candidates threatened the timely completion of the schedule intense protocol. Although one study suggests that SBO at spinal level S1 has a higher incidence of posterior disc herniation, the preponderance of clinical literature reports that spina bifida occulta is not a medical problem. The impact literature indicates that lumbosacral vertebral bodies fracture at 7.14 kN in static compression and 20 plus g during dynamic vertical impacts. In this paper, we examined the human data observed in ejection seat incidents, the rationale for excluding volunteers with single level SBO, and the path of axial load transmission through the lumbosacral spine. Based on the findings, we concluded that research volunteers with single level SBO are not at increased risk for injury, and recommended inclusion of these volunteers in future studies involving repeated axial impacts due to ride motion.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Chief, Science Support Center			22b. TELEPHONE (Include Area Code) (334) 255-6907		22c. OFFICE SYMBOL MCMR-UAX-SS

Contents

	Page
Introduction	3
Methods	3
Findings	4
Discussion	4
Conclusion	5
References	8

List of figures

Figure	Page
1. Path of axial load transmission	7

List of tables

Table	Page
1. Distribution of SBO in screened asymptomatic volunteers	4

This page left blank intentionally

INTRODUCTION

The nature of the various military occupational specialties dictate strict medical and physical standards. Some service members are exposed to repeated impacts with significant axial loads such as those in fast attack vehicles and high speed boats. Spinal dysraphic conditions other than spina bifida occulta (SBO) without overlying skin manifestations at one vertebral level are disqualifying for military duties (DOD Directive 6130.3). The military routinely does not screen for SBO radiographically.

Spina bifida occulta is a developmental abnormality that consists of a small interruption in continuity of the posterior arch with no neurologic deficit (15). The spinous process may be absent or is composed of two rudimentary processes from the laminae meeting, but not fused in the midline. The intervening space is occupied by a fibrous membrane. SBO, without cutaneous manifestations, is rarely associated with any clinical consequences (13). However, one study suggests that SBO at spinal level S1 has a higher incidence of posterior disc herniation (2). SBO is usually an incidental finding on radiographs of normal people (4). Single level SBO occurs commonly at spinal level S1, followed by L5, then in the cervical area (8). Surveys of normal, healthy adults show a prevalence between 18-34% (3,4,8,12,14). According to a 1946 study (7), the prevalence in the military population (mean age 26.6) is 36%.

The medical effects to single, high + Gz (axial) forces are well known in the ejection seat environment (10). However, the health and performance effects of repeated impact exposure are unknown but currently under study. A recent US Army Aeromedical Research Laboratory (USAARL) ride motion study excluded volunteer research subject candidates with SBO because of a perceived increased risk for injury. Disqualifying a significant percentage of screened research subject candidates threatened the timely completion of the schedule intense ride motion protocol. This paper examines the rationale for excluding research subject volunteers with SBO at one vertebral level and proposes a new guideline.

METHODS

Seventy nine male soldier volunteers, ages 18-40, were screened for exposure to repeated mechanical impacts up to 4 g at 5-20 Hz, and for durations up to 7 hours for one day and up to 4 hours on five consecutive days. Medical screening included a complete history, focused physical exam, and a standard anteroposterior and lateral lumbosacral series. The soldiers were disqualified from the participation in the study if their history indicated chronic symptoms, recurrent treatment, or surgical procedures for pathologic back conditions. A history of episodic mechanical low back pain was not a disqualifying criterion. They were disqualified if their physical exam revealed scoliosis, lower extremity neuropathy, cutaneous manifestations of SBO, or acute tenderness to palpation of the paravertebral musculature. Any radiographic variant or abnormality, as read by the board-certified radiologist of the adjacent accredited military medical facility, was exclusionary.

Female soldiers were not included in the protocol for two reasons. At the time the protocol was designed, female soldiers were not assigned to units that had the fast attack vehicles or high speed boats. Hence, female soldiers were not exposed to the impact forces in question. Also, choosing only male research subjects eliminated physiologic data variation between males and females.

FINDINGS

The mean age for the volunteer candidate population was 26.8. Forty-seven percent (37/79) of the male volunteer candidates were officers and 53% (42/79) were enlisted. Eighty-four percent (66/79) of the volunteer candidates were white and 16% (13/79) were nonwhite. All had a negative history of a chronic back problems and were asymptomatic for back pain.

Of the 17 medically disqualified from the protocol, one had just recovered from an acute episode of mechanical low back pain, one had spondylolysis, one had degenerative joint disease, and one had a pars defect. None of the remaining disqualified subjects had dimpling or a hairy patch over the lumbosacral spine. Thirteen had SBO at either the L5 or S1 level on radiographic examination. Table 1 compares those screened volunteers who had SBO to those who did not by age, race and rank.

Table I.
Distribution of SBO in screened asymptomatic volunteers.

		SBO	No SBO
N		13	66
AGE (mean)		27.7	26.6
RACE	Caucasian	12	54
	Noncaucasian	1	12
RANK	Officer	6	31
	Enlisted	7	35

There was no significant difference in the mean age for those who had SBO and those without SBO (T test=.7711, p=.4430). Univariate Logistic Regression model showed no association of SBO with race (Caucasian=1, OR=2.67, CI_{0.95}=0.316,22.5) and rank (officer=1, OR=0.968, CI_{0.95}=0.294,3.19).

DISCUSSION

The 16.5% prevalence of SBO in our population of male soldiers was consistent with the 23% found in the normal, asymptomatic U.S. population (6), 22% found in an emergency

department's outpatient population whose symptoms were completely unrelated (3), 15.2% found in a London historic population (13), 15.7% found in a contemporary London population (13), and 17% of a referred back pain population (2).

Based on the automotive impact literature, the compressive force required to fracture an isolated cadaver lumbar vertebra is 7.14 kN, the thoracic vertebra is 6.30 kN, and cervical vertebra is 4.09 kN (12). Assuming a 100-pound mass loads the spine (upper torso), these forces are equivalent to 16.1 G, 14.2 G and 9.2 G, respectively. The high performance jet community put a 20 G exposure limit on ejection seats (5). Single impact (ejection seat) primate studies showed fractures that clustered around C7 and T10 (10) and observational human ejection seat incidents showed clustering around T12/L1 (1). This suggests that the spinal area at risk is not the lumbosacral area and that the lumbosacral area can be subjected to higher loads than the cervicothoracic areas.

In normal backs, it is unknown if a significant amount of axial loading is transmitted through the pars interarticularis at L5, or whether the vertebral centrum bears the full load (11). Symmetrically loaded primate spines showed primarily comminuted, burst vertebral centrum fractures, while asymmetric loading showed wedge fractures and significant derangements of the articular facets (10). The lumbar articular facets are nonweight bearing and their involvement in force transmission is dependent on the rotational angle of the pelvis and the resiliency of the intervertebral disks (11). In a seated individual whose pelvis is rotated forward, the path of force transmission is through the anterior elements of the lumbosacral spine as shown in figure 1.

The U.S. Air Force had significant medical concerns for those who had back conditions that were previously quiescent and did not produce sufficient symptoms until exposure to the high G environment (9). These conditions included spondylolysis, spondylolisthesis, Klippel-Feil syndrome, Scheuermann's disease, spondylolysis deformans, and vertebral body fractures. It is believed that any structure with less than optimal strength is at increased risk for injury, however, biomechanical studies have not been done to correlate vertebral column abnormalities to spinal kinematics and strength (11). In the case of SBO at one vertebral level, it is reasonable that the axial force can be transmitted through the intact ipsilateral pars interarticularis.

Some limitations remain when evaluating whether asymptomatic SBOs are at higher risk for injury than those with normal backs. The fracture loads stated are for single static compressive and dynamic vertical impacts. We do not know if there is a fatiguing cumulative effect of repeated impacts that would lower the fracture threshold.

CONCLUSION

Because the prevalence of SBO inhibited efforts to recruit research subjects, timely execution and completion of a ride motion research project was threatened. Furthermore, conclusions [results] from the ride motion study may not be valid because a representative military population was not chosen. Given the vertebral fracture load forces, spinal level at risk, path of force transmission, clinical insignificance of SBO, and low level G exposure of repeated impacts in surface ride motion,

we do not feel that normal, asymptomatic research volunteers with SBO at one vertebral level and no cutaneous manifestations, are at increased risk for personal injury. We recommend that others designing similar studies consider including these research candidates.




Figure 1. Path of axial load transmission primarily through the anterior element with minimal transmission through the pars interarticularis in a normal lumbosacral spine. SBO at L5 or S1 would not affect the ipsilateral transmission through the pars interarticularis.

REFERENCES

1. Auffret R, Delahaye RP. Spinal injury after ejection. AGARD Advisory Report No.72. 1975;1-7.
2. Avrahami E, Frishman E, Fridman Z, Azor M. Spina bifida occulta of S1 is not an innocent finding. Spine. 1994; 19(1):12-15.
3. Boone D, Parsons D, Lachman M, Sherwood T. Spina bifida occulta: lesion or anomaly? Clin Radiol. 1985; 36:159-161.
4. D'Ambrosia RD. Musculoskeletal disorders: regional examination and differential diagnosis. Toronto: JP Lippincott Company. 1977; 311.
5. Ernsting J, King P. Aviation Medicine, 2nd ed; Boston: Butterworths. 1988; 169.
6. Fidas A et al. Prevalence and patterns of spina bifida occulta in 2707 normal adults. Clin Radiol. 1987; 38:537-542.
7. Friedman M, Fischer F, VanDemark R. Lumbo sacral roentgenograms of one hundred soldiers. American Journal of Radiology. 1946; 55(3):292-298.
8. Jinkins JR. Congenital anomalies of the lumbosacral spine. Neuroimaging Clin N Am: The lumbosacral spine. 1993; 3(3):429-430.
9. Kazarian LE, Belk WF. Flight physical standards of the 1980's: spinal column considerations. Wright-Patterson AFB, OH: Armstrong Aerospace Medical Research Laboratory. 1979. Technical Report 79-74.
10. Kazarian LE, Boyd D, von Gierke H. The dynamic biomedical nature of spinal fractures and articular facet derangement. Linear accelerations of impact type. In: AGARD Conference Proceedings No. CP-88-71. 1988:19.1-19.25.
11. Kazarian LE. Injuries to the human spinal column: biodynamics and injury classification. Wright-Patterson AFB, OH: Armstrong Aerospace Medical Research Laboratory. Technical Report 80-53, 1980.
12. Melvin JW, Weber K. Review of biomechanical impact response and injury in the automotive environment. U.S. Department of Transportation, 1986;
13. Salter RB. Textbook of disorders and injuries of the musculoskeletal system, 2nd ed. Baltimore: Williams and Wilkins. 1984; 133.

14. Saluja PG. The incidence of spina bifida occulta in a historic and modern London population. *J Anat* 1988; 158:91-93.
15. Turek SL. *Orthopedics: principles and their applications*, 4th ed. New York: JP Lippincott Company. 1984; 1552-1555.